Chaffinch (Fringilla coelebs)RBCommon throughout thesite.

Siskin (*Carduelis spinus*) W Occasionally seen with other finches in flocks during the winter months.

Greenfinch (*Carduelis chloris*) RB Most often seen in the hedgerows and shaws during spring/summer and with flocks of finches during the winter.

Goldfinch (*Carduelis carduelis*) RB Most often seen in the hedgerows and shaws during spring/summer and with flocks of finches during the winter.

Bullfinch (Pyrrhula pyrrhula)RBSmall numbers seen onwoodland perifery and in shaws/scrub.

Lesser Redpoll (*Carduelis cabaret*) W Large flock (140) present in January - March, single bird present in December.

Linnet (Carduelis cannabina)RBPair seen on Hitherfield in June. Pair nested at Harholt Plantation June.

Yellowhammer (*Emberiza citrinella*) R Single, singing male at Stagbury 'B' in May.

Ring-necked Parakeet (*Psittacula krameri*)RBSmall butnoisy colony present in the Nuttery.Small but

All photographs © Andy Keay 2011.



Juvenile Whitethroat

Key to status RESIDENT [R] BREEDING [B] SUMMER BREEDING VISITOR [S] WINTER VISITOR [W] VISITOR, ANY SEASON [V]

Red Kite (*Milvus milvus*) V Occasional visitor, usually over the grasslands in the Chipstead Valley.

Sparrowhawk (*Accipiter nisus*) R Although likely to be a breeding species, no nest was discovered in 2011. Frequently seen in and over Banstead Wood.

Buzzard (*Buteo buteo*) R Nested in an adjacent woodland, but not on the site. Rare twenty years ago, they are now seen increasingly and seem likely to become a breeding species.

Kestrel (*Falco tinnunculus*) RB Nested in the shaw between Harholt Plantation and Hill Wood Field. Frequently seen over the Downs.

Hobby (*Falco subbuteo*) S Nested in Banstead Wood producing two young. Possibly under threat due to substantial decrease in its main prey (Swallows, swifts and martins).

Pheasant (*Phasianus colchicus*) R Usually seen and heard along the western boundary of Banstead Wood where it may have bred.

Lapwing (*Vanellus vanellus*) W Seen over grassland areas on two occasions during the winter months.

Woodcock (*Scolopax rusticola*) R Two birds seen in February. Possible breeding species? Very prone to increasing disturbance (dogs, etc.)

Woodpigeon (Columba palumbus) RB Common breeding species.

Goldcrest (*Regulus regulus*) RB Small numbers particularly seen on conifers or accompanying flocks of tits in Banstead Wood.

Blue Tit (Parus caeruleus)	RB	Common throughout	
Great Tit (Parus major)	RB	Common throughout	
Coal Tit (Parus ater)RBGoldcrests on conifers in Banstea		mall numbers, often with	
Long-tailed Tit (<i>Aegithalos caud</i> heard, particularly in the Fames R	,	1 1	
Nuthatch (Sitta europaea)RBProbably increasing innumbers. Found throughout the woodland and shaws.			
Treecreeper (<i>Certhia familiaris</i>) throughout the woodlands.	RB	Small numbers distributed	
Starling (<i>Sturnus vulgaris</i>) serious decline. Rarely seen on th	R e site!	Resident species suffering	
Jay (<i>Garrulus glandarius</i>) heard although usually a shy bird.	RB	Frequently seen and	
Magpie (<i>Pica pica</i>) increasing throughout.	RB	Common and	
Jackdaw (Corvus monedula)RBLarge colony in theChipstead Valley, frequently feeding on the Downs.			
Raven (Corvus corax)RA single bird beingmobbed by Carrion Crows is probably attributable to this species. Itwas seen over the Downs in May.			
Carrion Crow (<i>Corvus corone</i>) site.	RB	Common throughout the	

House Sparrow (*Passer domesticus*) RB Serious decline in numbers and now only regularly seen around the Rambler's Rest.

Garden Warbler (Sylvia borin)	S	Male(s) heard singing in
Harholt Plantation.		

Lesser Whitethroat (Sylvia curruca)SMale singing atStagbury fieldS

Whitethroat (*Sylvia communis*) S Males singing at Stagbury field, Great Parson's field and Harholt Plantation.

Blackcap (*Sylvia atricapilla*) R Several males singing in scrubby areas in Banstead Wood and adjacent shaws/scrub. Increasingly over-wintering.



Willow Warbler (*Phylloscopus trochilus*) at Harholt Plantation.

Chiffchaff (*Psylloscopus collybita*) S F scrub areas and the perifery of Banstead Wood.

Present mostly

S

Male Blackcap singing

Present in most shaws,



Stock Dove (*Columba oenas*) R Infrequently seen and heard in Banstead Wood. Probable breeding species but not yet confirmed.

Collared Dove (*Streptopelia decaocto*) R Occasionally seen in the vicinity of the Rambler's Rest.

Tawny Owl (Strix aluco)RBOften heard butinfrequently seen in Banstead Wood and its immediate environs.

Little Owl (Athene noctua)RPossibly heard, but notyet seen in Banstead Wood.R

Swift (Apus apus)SInfrequently seen orheard over the site in 2011.

Green Woodpecker (*Picus viridis*) RB Maintaining good size population. Most often seen on Hither Field, Hill Wood field and Harholt Plantation.

Great Spotted Woodpecker (*Dendrocopus major*) RB Seen and heard freqently, mostly in Banstead Wood.

Lesser Spotted Woodpecker (*Dendrocopus minor*) R A single bird seen on the perifery of Banstead Wood in June. Possibly suffering from the decrease in food supply (invertebrates) due to weather conditions?

Skylark (*Alauda arvensis*) RB Fewer breeding birds on Hither field this year, probably due to the increasing scrub. No juvenile birds seen, so likely that the nesting birds failed to raise any young.

Swallow (*Hirundo rustica*) S Few of this migratory species seen over the site during the summer - due to decrease in invertebrates?

House Martin (*Delichon urbica*) S Very small numbers seen over the site during the summer - due to decrease in numbers of invertebrates?

Meadow Pipit (*Anthus pratensis*) R A species new to the site. Small flock seen on Hither field during September and October.

Pied Wagtail (Motacilla alba)RSeen infrequently atPark Farm depot and Ramber's Rest.

Wren (Troglodytes troglodytes)RBCommon breedingspecies in Banstead Wood, infrequent over the rest of the site.

Dunnock (*Prunella modularis*) RB Common breeding species on the perifery of Banstead Wood and hedgerows over the site.

Robin (Erithacus rubecula)RBCommon breedingspecies in Banstead Wood and hedgerows over the site.

Whinchat (Saxicola rubetra)SSingle male seenon Hither field, present for several days in late August.

Stonechat (Saxicola torquata)RSmall flock (3)seen on Hither field, present for one day in September.

Wheatear (Oenanthe oenanthe)SSingle bird seen onHither field in April.S

Blackbird (Turdus merula)RBSeveral breedingpairs in Banstead Wood and also in shaws/scrub on the Downs.

Fieldfare (Turdus pilaris)WModerate numbers,usually in company with Redwings, during January - March. Seen overall of the site, particularly close to scrub bearing berries!

Redwing (Turdus iliacus)WHigh numbers duringJanuary - March.V

Song Thrush (Turdus philomelos)RBSeveral pairs inBanstead Wood, rarely seen elsewhere on the site.

Mistle Thrush (Turdus viscivorus)RBAt least 3 pairs inBanstead Wood, with another singing male in the Nuttery.